

**Letter to the Editor****\*Corresponding author****Mehtap Honca, MD**Associate Professor  
Department of Anesthesiology and  
Reanimation  
Kecioren Training and Research  
Hospital

Ankara, Turkey

Tel. +90 312 3569000

Fax: +90 312 3569002

E-mail: [mehtaphonca@hotmail.com](mailto:mehtaphonca@hotmail.com)**Volume 1 : Issue 1****Article Ref. #: 1000AOJ1102****Article History****Received:** January 21<sup>st</sup>, 2016**Accepted:** January 25<sup>th</sup>, 2016**Published:** January 25<sup>th</sup>, 2016**Citation**Honca M. Pediatric emergence agita-  
tion. *Anesthesiol Open J.* 2016; 1(1):  
2-3. doi: [10.17140/AOJ-1-102](https://doi.org/10.17140/AOJ-1-102)**Copyright**©2016 Honca M. This is an open  
access article distributed under the  
Creative Commons Attribution 4.0  
International License (CC BY 4.0),  
which permits unrestricted use,  
distribution, and reproduction in  
any medium, provided the original  
work is properly cited.**Pediatric Emergence Agitation****Mehtap Honca, MD\****Department of Anesthesiology and Reanimation, Kecioren Training and Research Hospital,  
Ankara, Turkey*

Emergence Agitation (EA) is still remaining as a major problem during the early stage of recovery from general anesthesia in children. EA was first reported in 1960's and it has been considered as a mental disturbance during recovery from general anesthesia which consists of hallucinations, delusions and confusion manifested by moaning, restlessness, involuntary physical activity and thrashing about in the bed.<sup>1</sup> The incidence of EA is variable and can reach to 80%.<sup>2</sup>

Pre-school aged children, preoperative anxiety, inadequate pain control, type of the surgery (especially ophthalmological and otorhinolaryngological), anesthesia method are defined as the risk factors for EA in children.<sup>3-7</sup> Sevoflurane and desflurane are the preferred inhalational anesthetic agents for the induction of anesthesia in children. However, these agents have been reported to increase the ratio of EA because of their low blood/gas partition coefficients.<sup>8</sup> Although emerge times from propofol and sevoflurane are similar, EA is frequently observed in sevoflurane based anesthesia. In recent studies elevation of intraserebral glucose and lactic acid concentrations were observed with Magnetic Resonance Spectroscopy (MRS) with the use of sevoflurane.<sup>9</sup> Also epileptogenic activity of sevoflurane has been reported.<sup>10</sup> However the exact mechanism of EA has not been understood yet.

Different anesthetic agents such as opioids, midazolam, ketamine, alpha-2 agonist sedatives and nonsteroidal anti-inflammatory drugs have been used to prevent EA. Low dose ketamine added to propofol was found effective in the prevention of EA in children with a history of EA with propofol Total intravenous anaesthesia (TIVA).<sup>11</sup> Hadi et al Found that ketodex (lowdoseketamine 0,15 mg/kg followed by dexmedetomidine 0,3 µ/kg IV) reduced the incidence and severity of EA in children undergoing adenotonsillectomy following sevoflurane based anesthesia and provided smooth extubation.<sup>12</sup> µ-opioid agonists were found effective in decreasing the incidence of EA under sevoflurane anesthesia but postoperativenausea and vomiting were increased.<sup>13</sup> In a study by Fang et al, the efficacy of midazolam, dexmedetomidine, ketamine, fentanyl, and propofol were compared for the prevention of sevoflurane-related EA in children with placebo. They found that all of these agents have decreased the incidence of EA but dexmedetomidine was considered the most effective agent in their study.<sup>14</sup> In a study by Rosen et al, 42% of the anesthesiologists declared that EA was a significant problem at their institution and propofol was found the most common anesthetic agent used to prevent and to treat EA compared with the other medications.<sup>15</sup>

EA is a major problem in the postoperative period that may cause physical harm to the patient and can be disturbing to the parents and nursing staff in there recovery room. There is not defined spesific treatment and optimal anesthetic technique in current practice. For this reason further studies are needed to understand the mechanism of EA and improve the treatment.

**REFERENCES**

1. Wilson TA, Graves SA. Pediatric considerations in a general postanesthesia care unit. *J Post Anesth Nurs.* 1990; 5: 16-24.
2. Dahmani S, Mantz J, Veyckemans F. Case scenario: severe emergence agitation after myringotomy in a 3-yr-old child. *Anesthesiology.* 2012; 117: 399-406. doi: [10.1097/ALN](https://doi.org/10.1097/ALN).

0b013e31825fb069

3. Aono J, Ueda W, Mamiya K, Takimoto E, Manabe M. Greater incidence of delirium during recovery from sevoflurane anesthesia in preschool boys. *Anesthesiology*. 1997; 87: 1298-1300.
4. Kain ZN, Caldwell-Andrews AA, Maranets I, et al. Preoperative anxiety and emergence delirium and postoperative maladaptive behaviors. *Anesth Analg*. 2004; 99: 1648-1654.
5. Aouad MT, Kanazi GE, Siddik-Sayyid SM, Gerges FJ, Rizk LB, Baraka AS. Preoperative caudal block prevents emergence agitation in children followings evoflurane anesthesia. *Acta Anaesthesiol Scand*. 2005; 49: 300-304. doi: [10.1111/j.1399-6576.2005.00642.x](https://doi.org/10.1111/j.1399-6576.2005.00642.x)
6. Dahmani S, Stany I, Brasher C, et al. Pharmacological prevention of sevoflurane-and desflurane-related emergence agitation in children: a meta-analysis of published studies. *Br J Anaesth*. 2010; 104: 216-223. doi: [10.1093/bja/aep376](https://doi.org/10.1093/bja/aep376)
7. Voepel-Lewis T, Malviya S, Tait AR. A prospective cohort study of emergence agitation in the pediatric post anesthesia care unit. *Anesth Analg*. 2003; 96: 1625-1630. doi: [10.1213/01.ANE.0000062522.21048.61](https://doi.org/10.1213/01.ANE.0000062522.21048.61)
8. Welborn LG, Hannallah RS, Norden JM, Ruttimann UE, Callan CM. Comparison of emergence and recovery characteristics of sevoflurane, desflurane and halothane in pediatric ambulatory patients. *Anesth Analg*. 1996; 83: 917-920.
9. Jacob Z, Li H, Makaryus R, et al. Metabolomic profiling of children's brains undergoing general anesthesia with sevoflurane and propofol. *Anesthesiology*. 2012; 117: 1062-1071. doi: [10.1097/ALN.0b013e31826be417](https://doi.org/10.1097/ALN.0b013e31826be417)
10. Vakkuri AP, Seitsonen ER, Jäntti VH, et al. A rapid increase in the inspired concentration of desflurane is not associated with the pileptiformence phalogram. *Anesth Analg*. 2005; 101: 396-400.
11. Anghelescu DL, Rakes LC, Shearer JR, Bikhazi GB. Prevention of emerge agitation in seven children receiving low dose ketamine and propofol total intravenous anesthesia. *AANA J*. 2011; 79(3): 238-242.
12. Hadi SM, Saleh AJ, Tang YZ, Daoud A, Mei X, Ouyang W. The effect of KETODEX on the incidence and severity of emergence agitation in children undergoing adenotonsillectomy using sevoflurane based-anesthesia. *Int J Pediatr Otorhinolaryngol*. 2015; 79(5): 671-676. doi: [10.1016/j.ijporl.2015.02.012](https://doi.org/10.1016/j.ijporl.2015.02.012)
13. Tan Y, Shi Y, Ding H, Kong X, Zhou H, Tian J.  $\mu$ -Opioidagonists for preventing emergence agitation under sevoflurane anesthesia in children: a meta-analysis of randomized controlled trials. *Paediatr Anaesth*. 2016; 26(2): 139-150. doi: [10.1111/pan.12815](https://doi.org/10.1111/pan.12815)
14. Fang XZ, Gao J, Ge YL, Zhou LJ, Zhang Y. Network meta-analysis on the efficacy of dexmedetomidine, midazolam, ketamine, propofol, and fentanyl for the prevention of sevoflurane-related emergence agitation in children. *Am J Ther*. 2015.
15. Rosen HD, Mervitz D, Cravero JP. Pediatric emergence delirium: Canadian pediatric anesthesiologists' experience. *Paediatr Anaesth*. 2016; 26(2): 207-212. doi: [10.1111/pan.12812](https://doi.org/10.1111/pan.12812)