

Mini Review

*Corresponding author

Marc Pignitter, PhD

Faculty of Chemistry
Department of Nutritional and
Physiological Chemistry
University of Vienna
Althanstrasse 14 2B524
Vienna 1090, Austria
Tel. +43 1 4277 70621
Fax: +43 1 4277 9706
E-mail: marc.pignitter@univie.ac.at

Volume 1 : Issue 4

Article Ref. #: 1000AFTNSOJ1113

Article History

Received: June 25th, 2015

Accepted: July 14th, 2015

Published: July 15th, 2015

Citation

Pignitter M, Somoza V. Impact of dietary oxidized lipids on energy metabolism. *Adv Food Technol Nutr Sci Open J.* 2015; 1(4): 76-81. doi: [10.17140/AFTNSOJ-1-113](https://doi.org/10.17140/AFTNSOJ-1-113)

Copyright

©2015 Pignitter M. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Impact of Dietary Oxidized Lipids on Energy Metabolism

Marc Pignitter* and Veronika Somoza

Department of Nutritional and Physiological Chemistry, University of Vienna, Althanstrasse 14 2B524, Vienna 1090, Austria

ABSTRACT

Consumption of dietary fat is known to influence metabolic rate and metabolic pathways. Dietary intake of unoxidized polyunsaturated fatty acids was shown to lead to an increased metabolic rate. Identification of the underlying mechanism revealed that modifications of the energy metabolism are associated with modifications of membrane lipid composition leading to the membrane pacemaker theory of metabolism. Mitochondrial membranes were shown to adapt their lipids to the dietary fat composition. Dietary fat is commonly prepared by applying heat treatment to increase palatability. Heat treatment of food lipids result in the formation of oxidized lipids. Intake of oxidized lipids might affect energy metabolism in a different way than their corresponding unoxidized lipids. However, scientific literature of the effects of individual oxidized lipids found in heat-treated dietary fats on the energy metabolism relevant for metabolic syndrome, diabetes and obesity research is scarce. This review comprises current knowledge of the impact of unoxidized and oxidized lipids on the energy metabolism.

KEYWORDS: Oxidized lipids; Energy metabolism; Membrane pacemaker theory of metabolism.

ABBREVIATIONS: ATP: Adenosine triphosphate; DHA: Docosahexaenoic acid; EPA: Eicosapentaenoic acid; GIT: Gastro Intestinal Tract; LDLR: Low Density Lipoprotein Receptor; PUFA: Polyunsaturated fatty acids; ROS: Reactive Oxygen Species; TAG: Triacylglycerol.

ORIGINS OF DIETARY OXIDIZED LIPIDS

Lipids are macronutrients which predominantly serve as constituents of all membranes, provide energy, and are involved in cellular signaling. The most abundant dietary lipids are Triacylglycerols (TAGs), comprising approximately 80-95% of all dietary lipids.¹ Other main dietary lipids are phospholipids and sterols. One of the most prominent representatives of sterols is cholesterol. Besides playing a key role in the physical characteristics of membranes, cholesterol is the precursor for steroid hormones and bile acids. Cholesterol is a monounsaturated lipid, which makes it prone to oxidation comparable to other mono- and poly-unsaturated fatty acyl chains in TAGs and phospholipids. The susceptibility of fatty acids to oxidation strongly depends on the degree of unsaturation. A high number of double bonds decreases the energy required for detachment of the bis-allylic hydrogen. While abstraction of the allylic hydrogen atom in oleic acid requires 322 kJ/mol, it only needs 171 kJ/mol in linoleic acid.² Once lipid oxidation is initiated, lipid radicals are rearranged to form conjugated diene radicals, which, in the presence of molecular oxygen, form peroxy radicals. By generating hydroperoxy lipids, autooxidation propagates. Fatty acid hydroperoxides can be further decomposed to volatile short-chain aldehydes, ketones or alcohols *via* scission of the carbon chain. Degradation of fatty acid hydroperoxides without scission of the carbon chain leads to the formation of triacylglycerides with keto, epoxy, hydroxyl and aldehyde groups, the so called oxidized monomers. Fatty acid hydroperoxides can also undergo condensation reactions resulting in the production of oxidized dimers and oligomers. Due to cyclization reactions and isomerizations cyclic fatty acid monomers and *trans* fatty acids could be identified as degradation products of fatty acid hydroperoxides. For cholesterol hydro-

peroxide, which is generated by the abstraction of allylic C-7 hydrogen, the most predominant decomposition products are 7-ketocholesterol, 25-hydroxycholesterol, 7 α -hydroxycholesterol, 7 β -hydroxycholesterol, cholesterol-5 α ,6 α -epoxide, cholesterol-5 β ,6 β -epoxide and cholesterol-3,5,6-triol.^{3,4} The amount of lipid oxidation products formed in foods depends on environmental factors, such as temperature, irradiation, oxygen availability, presence of (anti)oxidants, metals and enzymes (lipoxygenases).

AMOUNT OF DIETARY OXIDIZED LIPIDS

Considerable amounts of dietary oxidized lipids have been quantified in Western diet due to processing of food. The amount of cholesterol oxidation products varies from 0.1 $\mu\text{g/g}$ beef⁵ to 18.7 $\mu\text{g/g}$ mortadella⁶ in meat and can reach up to 33.6 $\mu\text{g/g}$ anchovies⁷ as in sea food. In butter, the content of oxysterols ranges from 13.7 to 27.3 $\mu\text{g/g}$.⁸ Formation of lipid hydroperoxides in corn oil heated at 100 °C for 36 h in the presence of air was calculated to be 243 mmol/L, while in the untreated oil solely 0.9 mmol/L lipid hydroperoxides could be detected.⁹ Thus, heating promotes lipid oxidation. Heating of safflower oil by frying potato chips seven times for 10 min with one hour storage at room temperature between each frying resulted in an approximately 15-fold increase of the peroxide value.¹⁰ Cholesterol oxidation products were also shown to be elevated in roasted salmon treated at 200 °C for 30 min yielding 7.38 $\mu\text{g/g}$ compared to fried samples treated at 180 °C for 3 min in olive oil yielding solely 2.98 $\mu\text{g/g}$.³ Besides elevated temperatures, cold fluorescent light was shown to induce lipid oxidation.¹¹ Recently, we could demonstrate an increase of the peroxide value by 1473 \pm 1.79% ($p\leq 0.001$) after household-representative storage of soybean oil in the presence of cold fluorescent light for 56 days.¹¹ During the household-representative storage of the study oil an increasing oxygen-containing headspace was considered to mimic consumer handling. Due to the multiple environmental factors determining the kind and amount of lipid oxidation products formed during food processing quantitative exposure of lipid oxidation products to humans is hard to generalize. However, under defined processing conditions the susceptibility of each lipid-containing food product to oxidation can be determined, leading to quantifiable amounts of ingested lipids.

ABSORPTION OF OXIDIZED LIPIDS

The absorption and metabolism of 1-¹⁴C-methyl linoleate hydroperoxide was studied in rats.¹² It could be shown that the labeled methyl linoleate hydroperoxide and its labeled decomposition products were chiefly recovered from the stomach (48.0%), the expired ¹⁴CO₂ (30.5%) and the small intestine (9.3%) 24 h after intubation of the labeled compound. Another study with rats, which received 17 μmol labeled linoleic acid hydroperoxide and 18 μmol unoxidized linoleic acid, confirmed the high recovery of approximately 65% of linoleic acid hydroperoxide in the gastric lumen immediately after ingestion.¹³ It could be shown that the decomposition products, linoleic acid hydroxide, epoxyketones, 9-oxononanoic acid and hexanal in-

creased several minutes after administration of the linoleic acid hydroperoxides, suggesting that linoleic acid hydroperoxide decomposed over time to these products in the gastric lumen. A small percentage of 15.4% of the ingested linoleic acid hydroperoxide and its decomposition products was recovered in the gastric tissue 30 min after administration. Hexanal was shown to enter the small intestine and be absorbed into the blood. To this end, Kanazawa and Ashida¹³ suggested that trilinoleoylglycerol hydroperoxides are cleaved in the stomach by gastric lipases to the free oxidized fatty acid, which are partly absorbed by the gastric tissue and partly decomposed to secondary reaction products. Subsequently, the decomposition products are partially absorbed by the intestine.

Cholesterol oxidation products were also reported to be absorbed in the intestines by different species. However, the degree of absorption in rats, rabbits and humans differed among the cholesterol oxidation products, with 7 β -hydroxycholesterol, cholesterol-5 α ,6 α -epoxide and 7-ketocholesterol being chiefly absorbed.¹⁴⁻¹⁶ After absorption of oxysterols in the upper intestinal tract, the oxysterols are transferred in the blood within chylomicrons. The chylomicrons carry TAGs to tissues. The activity of endothelial lipoprotein lipase leads to the formation of chylomicron remnants, which are rapidly cleared by the liver. Thus, lipid oxidation products have been shown to be bioavailable.

IMPACT OF (OXIDIZED) LIPIDS ON ENERGY METABOLISM

Once absorbed fatty acids can be stored as triglyceride in adipose tissue, transferred to extra-hepatic tissue within lipoproteins and used for energy supply in any tissue containing mitochondria with oxygen availability. As mitochondria are the site of β -oxidation of fatty acids and ATP production, it plays a key role in energy supply. Total energy expenditure at rest is measured as basal metabolic rate in humans and animals. Several animal studies reported a correlation between dietary fatty acid profile and the basal metabolic rate in animals.¹⁷⁻¹⁹ It could be shown that feeding omega-3 or omega-6 enriched diets to rats led to an increase of the metabolic rate compared to rats fed a saturated fat diet. Human studies confirmed that increasing PUFA content in the diet was associated with an elevated metabolic rate.^{20,21} In a human cross-over study six healthy volunteers received a control diet ad libitum for 3 weeks and after a break of 10-12 weeks the subjects were administered the same control diet where 6 g visible fat per day was replaced by 6 g fish oil per day for 3 weeks.²² Dietary intake of fish oil significantly reduced the body fat mass by -0.88 \pm 0.16 kg compared to the intake of visible fat which led to an alteration of the body fat mass by -0.3 \pm 0.34 kg. In addition, an increase of the rate of lipid oxidation to 1.06 \pm 0.17 $\text{mg}\times\text{kg}^{-1}\times\text{min}^{-1}$ was obtained when fish oil was consumed compared to the intake of the visible fat (0.87 \pm 0.13 $\text{mg}\times\text{kg}^{-1}\times\text{min}^{-1}$).

A recent study with LDLR^{-/-} mice fed a Western diet for 16 weeks revealed that liver metabolites associated with lipid and amino acid pathways were chiefly affected by the diet as

determined by a non-targeted metabolomic approach.²³ Supplementation of the Western diet with EPA or DHA reduced the Western diet-induced effects, whereby feeding the mice with a DHA-supplemented Western diet reversed the Western diet-induced effects more pronouncedly. Hepatic C₂₀₋₂₂ omega 3 fatty acids and their oxidation products were demonstrated to be enhanced after administration of DHA-supplemented Western diet, whereas monounsaturates, omega 6 fatty acids and its corresponding oxidation products were significantly decreased. Metabolomic analyses identified, for instance, 18-hydroxy-5Z, 8Z, 11Z, 14Z, 16E-eicosapentaenoic acid and 17, 18-dihydroxy-eicosa-5, 8, 11, 14-tetraenoic acid, two omega 3 fatty acids-derived oxidation products. DHA-supplemented diet was shown to affect the hepatic lipid metabolism, explaining the protective effect of DHA against Western-diet-induced nonalcoholic steatohepatitis in mice.²³

So, dietary fat has an impact on the energy metabolism by modifying the metabolic rate and the metabolic pathways. However, the impact of processed food-derived lipid oxidation products on energy metabolism has not yet been addressed, despite the fact that many foods undergo processes before consumption to increase palatability (Figure 1).




Figure 1: Proposed mechanism of the potential effects of oxidized lipids on energy metabolism after ingestion of heat-treated lipids.

The molecular mechanisms explaining the role of di-

etary (unoxidized) fat on energy metabolism have been under thorough investigations.²⁴⁻²⁷ The energy metabolism has been suggested to be associated with membrane lipid composition.²⁷ In the field of comparative biology it could be demonstrated that species with high metabolic rates (endotherms) have highly polyunsaturated membranes while ectotherms with low metabolic rates are linked to cellular membranes which consists of more monounsaturated fatty acid acyl chains.²⁷ This finding led to the development of the so called membrane pacemaker⁷ theory of metabolism.²⁷ In particular, membrane composition affects Na⁺/K⁺ antiporter activity, which accounts for 10-60% of the resting metabolic rate.²⁴⁻²⁶ The activity of the membrane-bound Na⁺/K⁺-ATPase was strongly correlated with the DHA content of the surrounding phospholipids.²⁸ It was suggested that a decrease in the degree of membrane lipid polyunsaturation might reduce energy-consuming processes such as the activity of ion transporters.²⁹ The membrane fatty acid composition might affect membrane-bound proteins, thereby modifying intracellular signaling. One of the integral membrane proteins, the glucose transporter 1, for instance, covers an area of approximately 17 molecules of a phosphatidylcholine bilayer consisting of saturated fatty acid chains.³⁰ Thus, high membrane fluidity is required for the insertion of the glucose transporter into the membrane. Membrane fluidity is primarily determined by the membrane composition. Unsaturated hydrocarbon tails cause a greater surface of the cross-section of the cylindrical hydrocarbon part of the phospholipid molecule compared to saturated tails. As a consequence, the interaction energy between the two unsaturated fatty acid chains is reduced, leading to an enhanced membrane fluidity.³⁰ The mechanism of fatty acid uptake was found to be similar to the mechanism of glucose uptake.³¹ The membrane-located fatty acid transporters were reported to regulate lipid metabolism. As for the activity of the glucose transporter, an impact of the membrane flexibility, and thus degree of membrane lipid polyunsaturation, on the activity of the fatty acid transporters might be conceivable. The impact of oxidized lipids on the fatty acid and glucose uptake and any correlation to the membrane flexibility has not yet been studied.

The extent to which dietary lipids are incorporated into cellular membrane has been investigated previously.³² The physiological conformer-regulator paradigm was applied to quantitate the incorporation of dietary lipids into the membrane, whereby the membrane lipids were plotted against the dietary lipids. Even though dietary lipid composition was changed this change could not be reflected in plasma membrane (average slope 0.07). Membranes are, thus, homeostatically regulated independent of the dietary fatty acids. However, a conforming response to dietary fat was obtained when the PUFA balance of the diet was below 10% of the membrane composition.³³ An average slope of the relationship between dietary fats and membrane lipids was determined to be 0.95 for membrane lipids from heart, liver, muscle, brain and red blood cells.

More specifically, mitochondrial membrane phospholipids were shown to conform to dietary fatty acids. Hepatic mitochondrial membrane lipid composition of rats fed a rapeseed

oil-rich diet for 11, 22 and 33 days was changed compared to the mitochondrial membrane of rats fed a standard diet.³⁴ The modified diet induced a decrease in the saturated to unsaturated molar ratio and an increased incorporation of oleic acid in the major mitochondrial tetra-acyl phospholipid, cardiolipin. It was reported that cardiolipin, which comprises 10-20% of total mitochondrial phospholipids, is essential for mitochondrial ATP formation.³⁵ Several studies showed that dietary lipids can modify mitochondrial respiration and ROS formation.³⁶⁻³⁹ Polyunsaturated fatty acids as well as lipid oxidation products are known to activate uncoupling proteins leading to proton leak across the inner mitochondrial membrane without using the electrochemical gradient for ATP production.^{40,41} Brookes, et al.⁴² demonstrated that membrane unsaturation was positively correlated with proton permeability and metabolic rate suggesting that mitochondrial fatty acid composition might affect mitochondrial inner membrane proteins. Battino, et al.⁴³ investigated the effect of feeding fried oil to rats on their liver mitochondrial respiratory proteins. Intake of fried extra virgin olive oil rich in polar lipid oxidation products enhanced the hydroperoxide and the thiobarbituric acid reactive substances contents of mitochondrial membranes. In addition, it induced a stimulatory effect on the cytochrome c oxidase activity and increased the cytochrome c+c1 and cytochrome a+a3 content compared to the administration of non-fried extra virgin olive oil.

CONCLUSION

The impact of dietary unoxidized fatty acids on the energy metabolism has been under thorough investigation. However, the bioenergetic effect of oxidized lipids still needs to be elucidated in mechanistic studies. So far, there is only little evidence that oxidized lipids might exert differential effects on mitochondrial respiratory chain. A systematic approach of the impact of lipid oxidation products from differently processed dietary fats on the bioenergetic pathways would be of great importance for the general public and especially for patients suffering from metabolic disorders.

CONFLICTS OF INTEREST

The authors have nothing to disclose.

REFERENCES

1. Hamam F. Specialty lipids in health and disease. *Food Nutr Sci.* 2013; 4: 63-70. doi: [10.4236/fns.2013.49A1011](https://doi.org/10.4236/fns.2013.49A1011)
2. Pignitter M, Somoza V. Critical evaluation of methods for the measurement of oxidative rancidity in vegetable oils. *J Food Drug Anal.* 2012; 20: 772-777. doi: [10.6227/jfda.2012200305](https://doi.org/10.6227/jfda.2012200305)
3. Echarte M, Zulet MA, Astiasaran I. Oxidation process affecting fatty acids and cholesterol in fried and roasted salmon. *J Agric Food Chem.* 2001; 49: 5662-5667. doi: [10.1021/Jf010199e](https://doi.org/10.1021/Jf010199e)
4. Conchillo A, Ansorena D, Astiasaran I. Intensity of lipid oxidation and formation of cholesterol oxidation products during frozen storage of raw and cooked chicken. *J Sci Food Agric.* 2005; 85: 141-146. doi: [10.1002/Jfsa.1969](https://doi.org/10.1002/Jfsa.1969)
5. Boselli E, Rodriguez-Estrada MT, Fedrizzi G, Caboni MF. Cholesterol photosensitized oxidation of beef meat under standard and modified atmosphere at retail conditions. *Meat Sci.* 2009; 81: 224-229. doi: [10.1016/j.meatsci.2008.07.023](https://doi.org/10.1016/j.meatsci.2008.07.023)
6. Novelli E, Zanardi E, Ghiretti GP, et al. Lipid and cholesterol oxidation in frozen stored pork, salame Milano and Mortadella. *Meat Sci.* 1998; 48: 29-40. doi: [10.1016/S0309-1740\(97\)00072-7](https://doi.org/10.1016/S0309-1740(97)00072-7)
7. Shozen K, Ohshima T, Ushio H, Takiguchi A, Koizumi C. Effects of antioxidants and packing on cholesterol oxidation in processed anchovy during storage. *Food Sci Technol-Lebensm-Wiss Technol.* 1997; 30: 2-8. doi: [10.1006/fstl.1996.0129](https://doi.org/10.1006/fstl.1996.0129)
8. Pie JE, Spahis K, Seillan C. Evaluation of oxidative-degradation of cholesterol in food and food ingredients - identification and quantification of cholesterol oxides. *J Agric Food Chem.* 1990; 38: 973-979. doi: [10.1021/Jf00094a012](https://doi.org/10.1021/Jf00094a012)
9. Udilova N, Jurek D, Marian B, Gille L, Schulte-Hermann R, Nohl H. Induction of lipid peroxidation in biomembranes by dietary oil components. *Food Chem Toxicol.* 2003; 41: 1481-1489. doi: [10.1016/S0278-6915\(03\)00164-9](https://doi.org/10.1016/S0278-6915(03)00164-9)
10. Khatoon S, Krishna AGG. Assessment of oxidation in heated safflower oil by physical, chemical and spectroscopic methods. *J Food Lipids.* 1998; 5: 247-267. doi: [10.1111/j.1745-4522.1998.tb00123.x](https://doi.org/10.1111/j.1745-4522.1998.tb00123.x)
11. Pignitter M, Stolze K, Gartner S, et al. Cold fluorescent light as major inducer of lipid oxidation in soybean oil stored at household conditions for eight weeks. *J Agric Food Chem.* 2014; 62: 2297-2305. doi: [10.1021/jf405736j](https://doi.org/10.1021/jf405736j)
12. Bergan JG, Draper HH. Absorption and metabolism of 1-14C-methyl linoleate hydroperoxide. *Lipids.* 1970; 5: 976-982. doi: [10.1007/BF02533200](https://doi.org/10.1007/BF02533200)
13. Kanazawa K, Ashida H. Dietary hydroperoxides of linoleic acid decompose to aldehydes in stomach before being absorbed into the body. *Biochim Biophys Acta.* 1998; 1393: 349-361. doi: [10.1016/S0005-2760\(98\)00089-7](https://doi.org/10.1016/S0005-2760(98)00089-7)
14. Osada K, Sasaki E, Sugano M. Lymphatic absorption of oxidized cholesterol in rats. *Lipids.* 1994; 29: 555-559. doi: [10.1007/BF02536627](https://doi.org/10.1007/BF02536627)
15. Linseisen J, Wolfram G. Absorption of cholesterol oxidation products from ordinary foodstuff in humans. *Ann Nutr Metab.* 1998; 42: 221-230. doi: [10.1159/000012737](https://doi.org/10.1159/000012737)

16. Vine DF, Mamo CL, Beilin LJ, Mori TA, Croft KD. Dietary oxysterols are incorporated in plasma triglyceride-rich lipoproteins, increase their susceptibility to oxidation and increase aortic cholesterol concentration of rabbits. *J Lipid Res.* 1998; 39: 1995-2004.
17. Shimomura Y, Tamura T, Suzuki M. Less body fat accumulation in rats fed a safflower oil diet than in rats fed a beef tallow diet. *J Nutr.* 1990; 120: 1291-1296.
18. Pan DA, Storlien LH. Dietary lipid profile is a determinant of tissue phospholipid fatty acid composition and rate of weight gain in rats. *J Nutr.* 1993; 123: 512-519.
19. Takeuchi H, Matsuo T, Tokuyama K, Shimomura Y, Suzuki M. Diet-induced thermogenesis is lower in rats fed a lard diet than in those fed a high oleic acid safflower oil diet, a safflower oil diet or a linseed oil diet. *J Nutr.* 1995; 125: 920-925.
20. Jones PJ, Schoeller DA. Polyunsaturated:saturated ratio of diet fat influences energy substrate utilization in the human. *Metabolism.* 1988; 37: 145-151. doi: [10.1016/S0026-0495\(98\)90009-9](https://doi.org/10.1016/S0026-0495(98)90009-9)
21. van Marken Lichtenbelt WD, Mensink RP, Westerterp KR. The effect of fat composition of the diet on energy metabolism. *Z Ernährungswiss.* 1997; 36: 303-305.
22. Couet C, Delarue J, Ritz P, Antoine JM, Lamière F. Effect of dietary fish oil on body fat mass and basal fat oxidation in healthy adults. *Int J Obes Relat Metab Disord.* 1997; 21: 637-643.
23. Depner CM, Traber MG, Bobe G, et al. A metabolomic analysis of omega-3 fatty acid-mediated attenuation of western diet-induced nonalcoholic steatohepatitis in LDLR^{-/-} mice. *PLoS One.* 2013; 8: e83756. doi: [10.1371/journal.pone.0083756](https://doi.org/10.1371/journal.pone.0083756)
24. Else PL, Wu BJ. What role for membranes in determining the higher sodium pump molecular activity of mammals compared to ectotherms? *J Comp Physiol B.* 1999; 169: 296-302. doi: [10.1007/s003600050224](https://doi.org/10.1007/s003600050224)
25. Hulbert AJ, Else PL. Basal metabolic rate: history, composition, regulation, and usefulness. *Physiol Biochem Zool.* 2004; 77: 869-876. doi: [10.1086/422768](https://doi.org/10.1086/422768)
26. Else PL, Wu BJ, Storlien LH, Hulbert AJ. Molecular activity of Na⁺,K⁺-ATPase relates to the packing of membrane lipids. *Ann N Y Acad Sci.* 2003; 986: 525-526. doi: [10.1111/j.1749-6632.2003.tb07240.x](https://doi.org/10.1111/j.1749-6632.2003.tb07240.x)
27. Hulbert AJ, Else PL. Membranes as possible pacemakers of metabolism. *J Theor Biol.* 1999; 199: 257-274. doi: [10.1006/jtbi.1999.0955](https://doi.org/10.1006/jtbi.1999.0955)
28. Turner N, Else PL, Hulbert AJ. Docosahexaenoic acid (DHA) content of membranes determines molecular activity of the sodium pump: Implications for disease states and metabolism. *Naturwiss.* 2003; 90: 521-523. doi: [10.1007/s00114-003-0470-z](https://doi.org/10.1007/s00114-003-0470-z)
29. Hulbert AJ, Turner N, Storlien LH, Else PL. Dietary fats and membrane function: implications for metabolism and disease. *Biol Rev Camb Philos Soc.* 2005; 80: 155-169. doi: [10.1017/S1464793104006578](https://doi.org/10.1017/S1464793104006578)
30. Weijers R. Membrane flexibility and exercise: A guide to Type 2 diabetes mellitus. *J Diabet Metab.* 2013; S10: 003. doi: [10.4172/2155-6156.S10-003](https://doi.org/10.4172/2155-6156.S10-003)
31. Glatz JFC, Luiken JJFP, Bonen A. Membrane fatty acid transporters as regulators of lipid metabolism: Implications for metabolic disease. *Physiol Rev.* 2010; 90: 367-417. doi: [10.1152/physrev.00003.2009](https://doi.org/10.1152/physrev.00003.2009)
32. Abbott SK, Else PL, Hulbert AJ. Membrane fatty acid composition of rat skeletal muscle is most responsive to the balance of dietary n-3 and n-6 PUFA. *Br J Nutr.* 2010; 103: 522-529. doi: [10.1017/S0007114509992133](https://doi.org/10.1017/S0007114509992133)
33. Abbott SK, Else PL, Atkins TA, Hulbert AJ. Fatty acid composition of membrane bilayers: Importance of diet polyunsaturated fat balance. *Biochim Biophys Acta.* 2012; 1818: 1309-1317. doi: [10.1016/j.bbamem.2012.01.011](https://doi.org/10.1016/j.bbamem.2012.01.011)
34. Monteiro JP, Pereira CV, Silva AM, et al. Rapeseed oil-rich diet alters hepatic mitochondrial membrane lipid composition and disrupts bioenergetics. *Arch Toxicol.* 2013; 87: 2151-2163. doi: [10.1007/s00204-013-1068-7](https://doi.org/10.1007/s00204-013-1068-7)
35. Schlame M, Towbin JA, Heerdt PM, Jehle R, DiMauro S, Blanck TJ. Deficiency of tetralinoleoyl-cardiolipin in Barth syndrome. *Ann Neurol.* 2002; 51: 634-637. doi: [10.1002/ana.10176](https://doi.org/10.1002/ana.10176)
36. Barzanti V, Battino M, Baracca A, et al. The effect of dietary lipid changes on the fatty acid composition and function of liver, heart and brain mitochondria in the rat at different ages. *Br J Nutr.* 1994; 71: 193-202. doi: [10.1079/BJN19940126](https://doi.org/10.1079/BJN19940126)
37. Clandinin MT, Field CJ, Hargreaves K, Morson L, Zsigmond E. Role of diet fat in subcellular structure and function. *Can J Physiol Pharmacol.* 1985; 63: 546-556.
38. Ramsey JJ, Harper ME, Humble SJ, et al. Influence of mitochondrial membrane fatty acid composition on proton leak and H₂O₂ production in liver. *Comp Biochem Physiol B Biochem Mol Biol.* 2005; 140: 99-108. doi: [10.1016/j.cbpc.2004.09.016](https://doi.org/10.1016/j.cbpc.2004.09.016)
39. Yamaoka S, Urade R, Kito M. Mitochondrial function in rats is affected by modification of membrane phospholipids with dietary sardine oil. *J Nutr.* 1988; 118: 290-296.

40. Beck V, Jaburek M, Demina T, et al. Polyunsaturated fatty acids activate human uncoupling proteins 1 and 2 in planar lipid bilayers. *FASEB J.* 2007; 21: 1137-1144. doi: [10.1096/fj.06-7489com](https://doi.org/10.1096/fj.06-7489com)

41. Murphy MP, Echtay KS, Blaikie FH, et al. Superoxide activates uncoupling proteins by generating carbon-centered radicals and initiating lipid peroxidation: Studies using a mitochondria-targeted spin trap derived from alpha-phenyl-N-tert-butyl nitron. *J Biol Chem.* 2003; 278: 48534-48545. doi: [10.1074/jbc.M308529200M308529200](https://doi.org/10.1074/jbc.M308529200M308529200)

42. Brookes PS, Buckingham JA, Tenreiro AM, Hulbert AJ, Brand MD. The proton permeability of the inner membrane of liver mitochondria from ectothermic and endothermic vertebrates and from obese rats: Correlations with standard metabolic rate and phospholipid fatty acid composition. *Comp Biochem Physiol B Biochem Mol Biol.* 1998; 119: 325-334. doi: [10.1016/S0305-0491\(97\)00357-X](https://doi.org/10.1016/S0305-0491(97)00357-X)

43. Battino M, Quiles JL, Huertas JR, et al. Feeding fried oil changes antioxidant and fatty acid pattern of rat and affects rat liver mitochondrial respiratory chain components. *J Bioenerg Biomembr.* 2002; 34: 127-134. doi: [10.1023/A:1015128009826](https://doi.org/10.1023/A:1015128009826)